

Anomalous diffusion in the presence of external forces: exact time-dependent solutions and entropy

Constantino Tsallis

Department of Chemistry, Baker Laboratory, and Materials Science Center

Cornell University, Ithaca, NY 14853-1301

and

Centro Brasileiro de Pesquisas Físicas

*Rua Xavier Sigaud 150, 22290-180 – Rio de Janeiro – RJ, Brazil **

Dirk Jan Bukman

Department of Chemistry, Baker Laboratory, Cornell University, Ithaca, NY 14853-1301

(February 1, 2008)

Abstract

The optimization of the usual entropy $S_1[p] = -\int du \, p(u) \ln p(u)$ under appropriate constraints is closely related to the Gaussian form of the *exact* time-dependent solution of the Fokker-Planck equation describing an important class of *normal* diffusions. We show here that the optimization of the generalized entropic form $S_q[p] = \{1 - \int du \, [p(u)]^q\}/(q-1)$ (with $q = 1 + \mu - \nu \in \mathcal{R}$) is closely related to the calculation of the *exact* time-dependent solutions of a generalized, nonlinear, Fokker Planck equation, namely $\frac{\partial}{\partial t} p^\mu = -\frac{\partial}{\partial x} [F(x)p^\mu] + D \frac{\partial^2}{\partial x^2} p^\nu$, associated with *anomalous* diffusion in the presence of the external force $F(x) = k_1 - k_2 x$. Consequently, paradigmatic types of normal ($q = 1$) and anomalous ($q \neq 1$) diffusions occurring in

*Permanent address (tsallis@cat.cbpf.br)

a great variety of physical situations become *unified* in a single picture.

05.20.-y; 05.40.+j; 05.60.+w; 66.10.Cb

Anomalous diffusion is intensively studied nowadays, both theoretically and experimentally. It is observed, for instance, in CTAB micelles dissolved in salted water [1], the analysis of heartbeat histograms in a healthy individual [2], chaotic transport in laminar fluid flow of a water-glycerol mixture in a rapidly rotating annulus [3], subrecoil laser cooling [4], particle chaotic dynamics along the stochastic web associated with a d=3 Hamiltonian flow with hexagonal symmetry in a plane [5], conservative motion in a d=2 periodic potential [6], transport of fluid in porous media (see [7] and references therein), surface growth [7], and many other interesting physical systems. Its thermostatistical foundation (as it is known for *normal* diffusion) is naturally highly desirable and has, since long, been looked for (see, for instance, [8] and references therein). This goal was recently achieved by Alemany and Zanette [9] (see also [10]) for *Lévy-like* anomalous diffusion, in the context of a generalized, not necessarily *extensive* (additive), thermostatistics that has been recently proposed [11,12]. This thermostatistics (already applied to a considerable variety of physical systems [13] and optimization techniques [14]) is based upon the entropic form

$$S_q[p] \equiv \frac{1 - \int du [p(u)]^q}{q - 1} \quad (q \in \mathcal{R}) \quad (1)$$

which reduces, in the $q \rightarrow 1$ limit, to the standard Boltzmann-Gibbs entropy

$$S_1[p] \equiv - \int du p(u) \ln p(u) \quad (2)$$

We show here that an approach similar to that of [9] makes possible a quite satisfactory discussion of a sensibly different anomalous diffusion, namely of the *correlated* type, characterized by the following generalized, *nonlinear* Fokker-Planck equation:

$$\frac{\partial}{\partial t} [p(x, t)]^\mu = - \frac{\partial}{\partial x} \{ F(x) [p(x, t)]^\mu \} + D \frac{\partial^2}{\partial x^2} [p(x, t)]^\nu \quad (3)$$

where $(\mu, \nu) \in \mathcal{R}^2$, $D > 0$ is a (dimensionless) diffusion constant, $F(x) \equiv -dV(x)/dx$ is a (dimensionless) external force (*drift*) associated with the potential $V(x)$, and (x, t) is a (dimensionless) 1+1 space-time. Let us mention that, in variance with the correlated type we are focusing on here, the Lévy-like anomalous diffusion is associated with a *linear* equation, though in fractional derivatives [5].

We intend to consider here a specific (but very common) drift, namely characterized by $F(x) = k_1 - k_2 x$ ($k_1 \in \mathcal{R}$ and $k_2 \geq 0$; $k_2 = 0$ corresponds to the important case of constant external force, and $k_1 = 0$ corresponds to the so called Uhlenbeck-Ornstein process [15]). The particular case $\mu = \nu = 1$ corresponds to the standard Fokker-Planck equation, i.e., to *normal* diffusion. The particular case $F(x) = 0$ (no drift) has been considered by Spohn [7] for $\mu = 1$ and arbitrary ν (for instance, $\nu = 3$ satisfactorily describes a standard solid-on-solid model for surface growth), and has been extended by Duxbury [16] for arbitrary μ and ν . The case $(\mu, k_1) = (1, 0)$ has been considered by Plastino and Plastino [17]. Our present discussion recovers *all* of these as particular instances.

First, let us illustrate the procedure we intend to follow, by briefly reviewing normal diffusion ($\mu = \nu = 1$). We wish to optimize S_1 (given by Eq. (2)) with the constraints

$$\int du p(u) = 1 \quad , \quad (4)$$

$$\langle u - u_M \rangle_1 \equiv \int du (u - u_M) p(u) = 0 \quad (5)$$

and

$$\langle (u - u_M)^2 \rangle_1 \equiv \int du (u - u_M)^2 p(u) = \sigma^2 \quad , \quad (6)$$

u_M and σ being fixed *finite* real quantities. The optimization straightforwardly yields the solution

$$p_1(u) = \frac{e^{-\beta(u-u_M)^2}}{Z_1} \quad (7)$$

with

$$Z_1 = \int du e^{-\beta(u-u_M)^2} = (\pi/\beta)^{1/2} \quad (8)$$

where $\beta \equiv 1/T$ is the Lagrange parameter associated with the constraint (6) and satisfies $\beta = 1/(2\sigma^2)$.

On the basis of Eq. (7) we propose, for the $\mu = \nu = 1$ particular case of Eq. (3) (i.e., the standard Fokker-Planck equation), the *ansatz*

$$p_1(x, t) = \frac{e^{-\beta(t)[x-x_M(t)]^2}}{Z_1(t)} \quad (9)$$

with

$$\frac{\beta(t)}{\beta(0)} = \left[\frac{Z_1(0)}{Z_1(t)} \right]^\lambda \quad (10)$$

It follows straightforwardly that $\lambda = 2$,

$$\frac{\beta(t)}{\beta(0)} = \left[\frac{Z_1(0)}{Z_1(t)} \right]^2 = \frac{1}{\left[1 - \frac{2D\beta(0)}{k_2} \right] e^{-2k_2 t} + \frac{2D\beta(0)}{k_2}} \quad (11)$$

and

$$\frac{dx_M(t)}{dt} = k_1 - k_2 x_M(t) \quad (12)$$

hence

$$x_M(t) = \frac{k_1}{k_2} + \left[x_M(0) - \frac{k_1}{k_2} \right] e^{-k_2 t}. \quad (13)$$

To discuss the $k_2 = 0$ case we can use $e^{-2k_2 t} \sim 1 - 2k_2 t$, which implies $x_M(t) = x_M(0) + k_1 t$ and $1/\beta(t) = [1/\beta(0)] + 4Dt$, which, in the limit $t \rightarrow \infty$ (i.e., $t \gg 1/[4D\beta(0)]$), yields the familiar result $1/\beta(t) \sim 4Dt$. This result implies, in turn, the celebrated Einstein expression $\langle (x - x_M)^2 \rangle_1 \propto t$ for Brownian motion.

Let us now address the *general* (μ, ν) case. Following along the lines of Alemany and Zanette [9] (and the generic framework of the generalized thermostatics [11,12]) we now wish to optimize S_q (given by Eq. (1)). The constraints are Eq. (4),

$$\langle u - u_M \rangle_q \equiv \int du (u - u_M) [p(u)]^q = 0 \quad (14)$$

(which generalizes Eq. (5)) and

$$\langle (u - u_M)^2 \rangle_q \equiv \int (u - u_M)^2 [p(u)]^q = \sigma^2 \quad (15)$$

(which generalizes Eq. (6)). This is an appropriate moment for commenting that the reason for using $[p(u)]^q$ (instead of the familiar $p(u)$) in the constraints (14) and (15) is the (very

essential) fact that by doing so we *preserve* [12] *the Legendre structure of Thermodynamics* and (through the nonnegativity of C_q/q [18], where C_q denotes the specific heat) *guarantee thermodynamic stability*. Let us consistently stress that the constraint (14) is equivalent to $\langle u \rangle_q = \langle u_M \rangle_q$, but not to $\langle u \rangle_q = u_M$ (since, unless $q = 1$, $\langle u_M \rangle_q \neq u_M$). All these peculiarities are of course originated by the essential *nonextensivity* that the index q introduces in the theory. For example, if we have two *independent* systems A and B (i.e., $p_{A*B}(u_A, u_B) = p_A(u_A)p_B(u_B)$), we immediately verify that $S_q(A*B) = S_q(A) + S_q(B) + (1-q)S_q(A)S_q(B)$.

It is straightforward to see that the above described optimization of S_q yields

$$p_q(u) = \frac{[1 - \beta(1-q)(u - u_M)^2]^{\frac{1}{1-q}}}{Z_q} \quad (16)$$

with

$$Z_q = \int du [1 - \beta(1-q)(u - u_M)^2]^{\frac{1}{1-q}} \quad (17)$$

In the limit $q \rightarrow 1$, these equations reduce to Eqs. (7) and (8), respectively. The corresponding *ansatz* for solving Eq. (3) now is

$$p_q(x, t) = \frac{\{1 - \beta(t)(1-q)[x - x_M(t)]^2\}^{\frac{1}{1-q}}}{Z_q(t)} \quad (18)$$

with

$$\frac{\beta(t)}{\beta(0)} = \left[\frac{Z_q(0)}{Z_q(t)} \right]^\lambda. \quad (19)$$

(as before, $\beta(t)$ and $Z_q(t)$ are nothing but the scaling of space with time). A tedious (but straightforward) calculation yields $\lambda = 2\mu$, and $q = 1 + \mu - \nu$. An equation for $Z_q(t)$ is also found, namely

$$2\nu D\beta(0)[Z_q(0)]^{2\mu} - k_2[Z_q(t)]^{\mu+\nu} - \frac{\mu}{\mu + \nu} \frac{d[Z_q(t)]^{\mu+\nu}}{dt} = 0, \quad (20)$$

which can be solved by substituting $\tilde{Z}(t) = Z_q(t)^{\mu+\nu}$. The resulting solution is (*for all values of k_1*)

$$Z_q(t) = Z_q(0) \left[\left(1 - \frac{1}{K_2}\right) e^{-t/\tau} + \frac{1}{K_2} \right]^{\frac{1}{\mu+\nu}} \quad (21)$$

with

$$K_2 \equiv \frac{k_2}{2\nu D\beta(0)[Z_q(0)]^{\mu-\nu}} \quad (22)$$

and

$$\tau \equiv \frac{\mu}{k_2(\mu + \nu)} \quad (23)$$

(See Fig. 1). The function $x_M(t)$ is the same as in the case of normal diffusion, Eq. (13), since it only describes the motion of the average of the distribution $p_q(x, t)$, and does not depend on the way in which it spreads. $\beta(0)$ and $Z_q(0)$ are determined by the initial condition (i.e., by $p_q(x, 0)$).

For $k_2 = 0$, Eq. (21) becomes

$$Z_q(t) = \left\{ [Z_q(0)]^{\mu+\nu} + \frac{2\nu(\nu + \mu)D\beta(0)[Z_q(0)]^{2\mu}}{\mu} t \right\}^{\frac{1}{\mu+\nu}} \quad (24)$$

which, for $t \rightarrow \infty$, asymptotically recovers Duxbury's solution [16], namely $1/\beta(t) \propto [Z_q(t)]^{2\mu} \propto t^{\frac{2\mu}{\mu+\nu}}$. As we see, $\mu/\nu = 1$, > 1 and < 1 respectively imply that $[x(t) - x_M(t)]^2$ scales like t (*normal diffusion*), faster than t (*superdiffusion*) and slower than t (*subdiffusion*). The limits $\mu/\nu = 0$ and $\mu/\nu = \infty$ correspond to “no diffusion” and ballistic motion, respectively.

For $(\mu, k_1) = (1, 0)$, the present set of equations reduces to that of Plastino and Plastino [17].

Let us mention that the general solution given in Eq. (21) can be derived from that for $\mu = 1$ (and arbitrary k_1) by defining $\tilde{p}(x, t) = [p(x, t)]^\mu$, and $\tilde{\nu} = \nu/\mu$, as can easily be seen from Eq. (3).

Finally, by using Eq. (19) with $\lambda = 2\mu$, we can verify that

$$\int dx p_q(x, t) = [Z_q(t)/Z_q(0)]^{\mu-1} \int dx p_q(x, 0) . \quad (25)$$

Consequently, the norm ("total mass") is generically conserved for all times only if $\mu = 1$ ($\forall K_2$) or if $K_2 = 1$ ($\forall \mu$). For $0 \leq K_2 < 1$ (a common case), the norm monotonically *increases* (*decreases*) with time if $\mu > 1$ ($\mu < 1$). If $K_2 > 1$, it is the other way around.

Before ending let us mention that, *also* when t grows to infinity, the solutions we have found must be physically meaningful. This imposes $\mu/\nu > -1$. Indeed, if $k_2 \neq 0$, τ in Eq. (21) must be positive, which implies $\mu/\nu > -1$. Also, if $k_2 = 0$, x must scale with an *increasing* function of t ; hence, $\beta(t)$ must *decrease* with t , which implies (through Eqs. (19) and (24)) $2\mu/(\mu + \nu) > 0$, hence, the already mentioned restriction applies once again. The entire picture which emerges is indicated in Fig. 2 (we have not focused the $\mu < 0$ region because that would force us to discuss the stability of the solutions with respect to small departures, and this lies outside of the scope of the present work).

Summarizing, on *general* grounds, we have shown that thermostatics allowing for nonextensivity constitute a theoretical framework within which a rather nice *unification* of normal and *correlated* anomalous diffusions can be achieved. Both types of diffusions have been founded, on equal footing, on primary concepts of (appropriately generalized) Thermodynamics and Information Theory.

On *specific* grounds, we have obtained, for a generic linear force $F(x)$, the physically relevant *exact* (space, time)-dependent solutions of a considerably generalized Fokker-Planck equation, namely Eq. (3).

It is with pleasure that one of us (C.T.) acknowledges warm hospitality by B. Widom at the Baker Laboratory. This work was carried out in the research group of B. Widom, and was supported by the National Science Foundation and the Cornell University Materials Science Center.

REFERENCES

- [1] A. Ott, J.P. Bouchaud, D. Langevin and W. Urbach, Phys. Rev. Lett. **65**, 2201 (1990);
J.P. Bouchaud, A. Ott, D. Langevin and W. Urbach, J. Phys. II France **1**, 1465 (1991).
- [2] C.-K. Peng, J. Mietus, J.M. Hausdorff, S. Havlin, H.E. Stanley and A.L. Goldberger, Phys. Rev. Lett. **70**, 1343 (1993).
- [3] T.H. Solomon, E.R. Weeks and H.L. Swinney, Phys. Rev. Lett. **71**, 3975 (1993).
- [4] F. Bardou, J.P. Bouchaud, O. Emile, A. Aspect and C. Cohen-Tannoudji, Phys. Rev. Lett. **72**, 203 (1994).
- [5] G.M. Zaslavsky, D. Stevens and H. Weitzner, Phys. Rev. E **48**, 1683 (1993); G.M. Zaslavsky, Physica D **76**, 110 (1994) and Chaos **4**, 25 (1994) and references therein.
- [6] J. Klafter and G. Zumofen, Phys. Rev. E **49**, 4873 (1994).
- [7] H. Spohn, J. Phys. I France **3**, 69 (1993).
- [8] M.F. Shlesinger and B.D. Hughes, Physica A **109**, 597 (1981); E.W. Montroll and M.F. Shlesinger, J. Stat. Phys. **32**, 209 (1983); E.W. Montroll and M.F. Shlesinger, in "Non-equilibrium Phenomena II: from Stochastic to Hydrodynamics", eds. J.L. Lebowitz and E.W. Montroll (North-Holland, Amsterdam, 1984).
- [9] P.A. Alemany and D.H. Zanette, Phys. Rev. E **49**, 956 (1994).
- [10] C. Tsallis, A.M.C. Souza and R. Maynard, in "Lévy flights and related topics in Physics", eds. M.F. Shlesinger, G.M. Zaslavsky and U. Frisch (Springer, Berlin, 1995), p. 269; D.H. Zanette and P.A. Alemany, Phys. Rev. Lett. **75**, 366 (1995); C. Tsallis, S.V.F. Levy, A.M.C. Souza and R. Maynard, Phys. Rev. Lett. **75** (Oct/Nov 1995), in press.
- [11] C. Tsallis, J. Stat. Phys. **52**, 479 (1988).

- [12] E.M.F. Curado and C. Tsallis, J. Phys. A **24**, L69 (1991); Corrigenda: **24**, 3187 (1991) and **25**, 1019 (1992).
- [13] A.R. Plastino and A. Plastino, Phys. Lett. A **174**, 384 (1993); J.J. Aly, in "N-Body Problems and Gravitational Dynamics", Proc. of the Meeting held at Aussois-France (21-25 March 1993), eds. F. Combes and E. Athanassoula (Publications de l'Observatoire de Paris, 1993), p. 19; A.R. Plastino and A. Plastino, Phys. Lett. A **193**, 251 (1994); L.S. Lucena, L.R. da Silva and C. Tsallis, Phys. Rev. E **51**, 6247 (1995); P. Jund, S.G. Kim and C. Tsallis, Phys. Rev. B **52**, 50 (1995); C. Tsallis, F.C. Sá Barreto and E.D. Loh, Phys. Rev. E **52**, 1447 (1995); B.M. Boghosian, preprint (1995) [E-mail to chaodyn@xyz.lanl.gov with "get chaodyn/9505012" on subject line]; M. Portesi, A. Plastino and C. Tsallis, Phys Rev E **52**, 3317 (1995).
- [14] T.J.P. Penna, Phys. Rev. E **51**, R1 (1995) and Computer in Physics **9**, 341 (1995); D.A. Stariolo and C. Tsallis, Ann. Rev. Comp. Phys., vol. II, ed. D. Stauffer (World Scientific, Singapore, 1995), p. 343; K.C. Mundim and C. Tsallis, Int. J. Quantum Chem. (1996), in press.
- [15] G.E. Uhlenbeck and L.S. Ornstein, Phys. Rev. **36**, 823 (1930).
- [16] P.M. Duxbury, private communication (1994); see [13] of P. Jund, S.G. Kim and C. Tsallis, Phys. Rev. B **52**, 50 (1995), and [18] of C. Tsallis, F.C. Sá Barreto and E.D. Loh, Phys. Rev. E **52**, 1447 (1995).
- [17] A.R. Plastino and A. Plastino, Physica A (1995), in press.
- [18] E.P. da Silva, C. Tsallis and E.M.F. Curado, Physica A **199**, 137 (1993); Erratum: Physica A **203**, 160 (1994).

FIGURES

FIG. 1. The $\mu/\nu = 1/3$ example: (a) Time dependence of $\beta(0)/\beta(t) = [Z_q(t)/Z_q(0)]^{2\mu}$ for $Z_q(0) \neq 0$ and typical values of K_2 (indicated at the right of each curve). The curve for $K_2 = 0$ lies on the vertical axis. For $K_2 = 0.25, 0.5$, and 2 the asymptotic values for $t/\tau \rightarrow \infty$ are shown by the dashed lines. (τ is defined in Eq. (23).); (b) Time dependence of $\{\beta(0)[Z_q(0)]^{2\mu}\}/\beta(t) = [Z_q(t)]^{2\mu}$ for $Z_q(0) = 0$, $\beta(0)[Z_q(0)]^{2\mu} \neq 0$, and typical values of $K_2' \equiv k_2/\{2\nu D\beta(0)[Z_q(0)]^{2\mu}\}$ (indicated at the right of each curve). The curve for $K_2 = \infty$ coincides with the horizontal axis. All curves saturate at a finite value as $t \rightarrow \infty$, except that for $K_2' = 0$, which is proportional to $t^{2\mu/(\mu+\nu)}$ for all t .

FIG. 2. "Norm conservation" means that $N \equiv \int dx p_q(x, t)$ is time-invariant; "Norm creation" means that N monotonically increases (decreases) with time if $K_2 < 1$ ($K_2 > 1$); "Norm dissipation" means that N monotonically decreases (increases) with time if $K_2 < 1$ ($K_2 > 1$). "Normal diffusion", "Superdiffusion" and "Subdiffusion" refer to the fact that, for $k_2 = 0$, $(x - x_M)^2$ scales like t , faster than t and slower than t , respectively. The standard Fokker-Planck equation corresponds to $\mu = \nu = q = 1$. For the precise meaning of "unphysical", see the text. On the $\mu = 1$ line we have $q = 2 - \nu$; consequently, when ν varies from ∞ to -1 , q varies from $-\infty$ to 3 , which precisely is the interval within which Eq. (4) (and, consistently, $\int dx p_q(x, 0) = 1$) can be satisfied.





